arXiv:math/0305232v4 [math.MG] 26 Jun 2003

Mappings from $\mathbb{R}^n$ to $\mathbb{F}^n$ which preserve unit Euclidean distance, where $\mathbb{F}$ is a field of characteristic 0

Apoloniusz Tyszka

Summary. Let $\mathbb{F}$ be a commutative field of characteristic 0, $\varphi_n : \mathbb{F}^n \times \mathbb{F}^n \to \mathbb{F}$, $\varphi_n((x_1, ..., x_n), (y_1, ..., y_n)) = (x_1 - y_1)^2 + ... + (x_n - y_n)^2$. We say that $g : \mathbb{R}^n \to \mathbb{F}^n$ preserves distance $d \geq 0$ if for each $x, y \in \mathbb{R}^n$ $|x - y| = d$ implies $\varphi_n(g(x), g(y)) = d^2$. Let $f : \mathbb{R}^n \to \mathbb{F}^n$ preserve unit distance. We prove: **(1)** if $n \geq 2$, $x, y \in \mathbb{R}^n$ and $x \neq y$, then $\varphi_n(f(x), f(y)) \neq 0$, **(2)** if $A, B, C, D \in \mathbb{R}^2$, $r \in \mathbb{Q}$ and $\overrightarrow{CD} = r\overrightarrow{AB}$, then $\overrightarrow{f(C)f(D)} = r\overrightarrow{f(A)f(B)}$, **(3)** if $A, B, C, D \in \mathbb{R}^2$ and $\overrightarrow{AB}$ and $\overrightarrow{CD}$ are linearly dependent, then $\overrightarrow{f(A)f(B)}$ and $\overrightarrow{f(C)f(D)}$ are linearly dependent, **(4)** if $A, B, C, D \in \mathbb{R}^2$ and $\overrightarrow{AB}$ is perpendicular to $\overrightarrow{CD}$, then $\overrightarrow{f(A)f(B)}$ is perpendicular to $\overrightarrow{f(C)f(D)}$, **(5)** if $A, B, C, D \in \mathbb{R}^2$ and $|AB| = |CD|$, then $\varphi_2(f(A), f(B)) = \varphi_2(f(C), f(D))$. Let $A_n(\mathbb{F})$ denote the set of all positive numbers d such that any map $g : \mathbb{R}^n \to \mathbb{F}^n$ that preserves unit distance preserves also distance d. Let $D_n(\mathbb{F})$ denote the set of all positive numbers d with the property: if $x, y \in \mathbb{R}^n$ and $|x - y| = d$ then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that any map map $g : S_{xy} \to \mathbb{F}^n$ that preserves unit distance preserves also the distance between x and y. Obviously, $\{1\} \subseteq D_n(\mathbb{F}) \subseteq A_n(\mathbb{F})$. We prove: **(6)** $A_n(\mathbb{C}) \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$, **(7)** $\{d > 0 : d^2 \in \mathbb{Q}\} \subseteq D_2(\mathbb{F})$.

2000 Mathematics Subject Classification: 51K05, 51M05.

Key words and phrases: Beckman-Quarles theorem, Cayley-Menger determinant, endomorphism (automorphism) of the field of complex numbers, isometry, unit-distance preserving mapping, unit Euclidean distance.

Let $\mathbb{F}$ be a commutative field of characteristic 0, $\varphi_n : \mathbb{F}^n \times \mathbb{F}^n \to \mathbb{F}$, $\varphi_n((x_1, ..., x_n), (y_1, ..., y_n)) = (x_1 - y_1)^2 + ... + (x_n - y_n)^2$. We say that $f : \mathbb{R}^n \to \mathbb{F}^n$ preserves distance $d \geq 0$ if for each $x, y \in \mathbb{R}^n$ $|x - y| = d$ implies $\varphi_n(f(x), f(y)) = d^2$. In this paper we study unit-distance preserving mappings from $\mathbb{R}^n$ to $\mathbb{F}^n$.

By a *complex isometry* of $\mathbb{C}^n$ we understand any map $h : \mathbb{C}^n \to \mathbb{C}^n$ of the form

$$h(z_1, z_2, ..., z_n) = (z'_1, z'_2, ..., z'_n)$$

where

$$z'_j = a_{0j} + a_{1j}z_1 + a_{2j}z_2 + ... + a_{nj}z_n \quad (j = 1, 2, ..., n),$$

the coefficients a_{ij} are complex and the matrix $||a_{ij}||$ $(i, j = 1, 2, ..., n)$ is orthogonal i.e. satisfies the condition

$$\sum_{j=1}^{n} a_{\mu j} a_{\nu j} = \delta_{\nu}^{\mu} \quad (\mu, \nu = 1, 2, ..., n)$$

with Kronecker's delta. According to [6], $\varphi_n(x, y)$ is invariant under complex isometries i.e. for every complex isometry $h : \mathbb{C}^n \to \mathbb{C}^n$

$$(\diamond) \qquad \forall x, y \in \mathbb{C}^n \ \ \varphi_n(h(x), h(y)) = \varphi_n(x, y).$$

Conversely, if $h : \mathbb{C}^n \to \mathbb{C}^n$ satisfies $(\diamond)$ then h is a complex isometry; it follows from [4, proposition 1, page 21] by replacing $\mathbb{R}$ with $\mathbb{C}$ and $d(x, y)$ with $\varphi_n(x, y)$. Similarly, if $f : \mathbb{R}^n \to \mathbb{C}^n$ preserves all distances, then there exists a complex isometry $h : \mathbb{C}^n \to \mathbb{C}^n$ such that $f = h_{|\mathbb{R}^n}$.

By a field endomorphism of $\mathbb{C}$ we understand any map $g : \mathbb{C} \to \mathbb{C}$ satisfying:

$$\forall x, y \in \mathbb{C} \ \ g(x + y) = g(x) + g(y),$$
$$\forall x, y \in \mathbb{C} \ \ g(x \cdot y) = g(x) \cdot g(y),$$
$$g(0) = 0,$$
$$g(1) = 1.$$

Bijective endomorphisms are called automorphisms, for more information on field endomorphisms and automorphisms of $\mathbb{C}$ the reader is referred to [10], [11] and [16]. If r is a rational number, then $g(r) = r$ for any field endomorphism $g : \mathbb{C} \to \mathbb{C}$. Proposition 1 shows that only rational numbers r have this property:

Proposition 1 ([15]). If $r \in \mathbb{C}$ and r is not a rational number, then there exists a field automorphism $g : \mathbb{C} \to \mathbb{C}$ such that $g(r) \neq r$.

Proof. Note first that if E is any subfield of $\mathbb{C}$ and if g is an automorphism of E, then g can be extended to an automorphism of $\mathbb{C}$. This follows from [7, corollary 1, A.V.111] by taking $\Omega = \mathbb{C}$ and $K = \mathbb{Q}$. Now let $r \in \mathbb{C} \setminus \mathbb{Q}$. If r is algebraic over $\mathbb{Q}$, let E be the splitting field in $\mathbb{C}$ of the minimal polynomial μ of r over $\mathbb{Q}$ and let $r' \in E$ be any other root of μ. Then there exists an automorphism g of E that sends r to r', see for example [9, corollary 2, page 66]. If r is transcendental over $\mathbb{Q}$, let $E = \mathbb{Q}(r)$ and let $r' \in E$ be any other generator of E (e.g. $r' = 1/r$). Then there exists an automorphism g of E that sends r to r'. In each case, g can be extended to an automorphism of $\mathbb{C}$.

If $g : \mathbb{C} \to \mathbb{C}$ is a field endomorphism then $(g_{|\mathbb{R}}, ..., g_{|\mathbb{R}}) : \mathbb{R}^n \to \mathbb{C}^n$ preserves all distances $\sqrt{r}$ with rational $r \geq 0$. Indeed, if $(x_1 - y_1)^2 + ... + (x_n - y_n)^2 = (\sqrt{r})^2$ then

$$\begin{aligned}
\varphi_n((g_{|\mathbb{R}}, ..., g_{|\mathbb{R}})(x_1, ..., x_n), (g_{|\mathbb{R}}, ..., g_{|\mathbb{R}})(y_1, ..., y_n)) &= \\
\varphi_n((g(x_1), ..., g(x_n)), (g(y_1), ..., g(y_n))) &= \\
(g(x_1) - g(y_1))^2 + ... + (g(x_n) - g(y_n))^2 &= \\
(g(x_1 - y_1))^2 + ... + (g(x_n - y_n))^2 &= \\
g((x_1 - y_1)^2) + ... + g((x_n - y_n)^2) &= \\
g((x_1 - y_1)^2 + ... + (x_n - y_n)^2) &= g((\sqrt{r})^2) = g(r) = r = (\sqrt{r})^2.
\end{aligned}$$

Analogously, if $g : \mathbb{R} \to \mathbb{F}$ is a field homomorphism then $(g, ..., g) : \mathbb{R}^n \to \mathbb{F}^n$ preserves all distances $\sqrt{r}$ with rational $r \geq 0$.

Conjecture 1. Each unit-distance preserving mapping from $\mathbb{R}^n$ to $\mathbb{F}^n$ $(n \geq 2)$ has a form $I \circ (g, ..., g)$, where $g : \mathbb{R} \to \mathbb{F}$ is a field homomorphism and $I : \mathbb{F}^n \to \mathbb{F}^n$ is an affine mapping with orthogonal linear part.

Theorem 1 ([15]). If $x, y \in \mathbb{R}^n$ and $|x - y|^2$ is not a rational number, then there exists $f : \mathbb{R}^n \to \mathbb{C}^n$ that does not preserve the distance between x and y although f preserves all distances $\sqrt{r}$ with rational $r \geq 0$.

Proof. There exists an isometry $I : \mathbb{R}^n \to \mathbb{R}^n$ such that $I(x) = (0, 0, ..., 0)$ and $I(y) = (|x - y|, 0, ..., 0)$. By Proposition 1 there exists a field automorphism $g : \mathbb{C} \to \mathbb{C}$ such that $g(|x-y|^2) \neq |x-y|^2$. Thus $g(|x-y|) \neq |x-y|$ and $g(|x-y|) \neq -|x-y|$. Therefore $(g_{|\mathbb{R}}, ..., g_{|\mathbb{R}}) : \mathbb{R}^n \to \mathbb{C}^n$ does not preserve the distance between $(0, 0, ..., 0) \in \mathbb{R}^n$ and $(|x - y|, 0, ..., 0) \in \mathbb{R}^n$ although $(g_{|\mathbb{R}}, ..., g_{|\mathbb{R}})$ preserves all distances $\sqrt{r}$ with rational $r \geq 0$. Hence $f := (g_{|\mathbb{R}}, ..., g_{|\mathbb{R}}) \circ I : \mathbb{R}^n \to \mathbb{C}^n$ does not preserve the distance between x and y although f preserves all distances $\sqrt{r}$ with rational $r \geq 0$.

Let $A_n(\mathbb{F})$ denote the set of all positive numbers d such that any map $f : \mathbb{R}^n \to \mathbb{F}^n$ that preserves unit distance preserves also distance d. The classical Beckman-Quarles theorem states that each unit-distance preserving mapping from $\mathbb{R}^n$ to $\mathbb{R}^n$ $(n \geq 2)$ is an isometry, see [1]-[4] and [8]. It means that for each $n \geq 2$ $A_n(\mathbb{R}) = (0, \infty)$. By Theorem 1 $A_n(\mathbb{C}) \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$. Let $D_n(\mathbb{F})$ denote the set of all positive numbers d with the following property:

$(*)$ if $x, y \in \mathbb{R}^n$ and $|x-y| = d$ then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that any map $f : S_{xy} \to \mathbb{F}^n$ that preserves unit distance preserves also the distance between x and y.

Obviously, $\{1\} \subseteq D_n(\mathbb{F}) \subseteq A_n(\mathbb{F})$. From [13] and [14] follows that for each $n \geq 2$ $D_n(\mathbb{R})$ is equal to the set of positive algebraic numbers. The author proved in [15] that for each $n \geq 2$ $D_n(\mathbb{C})$ is a dense subset of $(0, \infty)$. The proof remains valid for $D_n(\mathbb{F})$.

Theorem 2. If $n \geq 2$ and $f : \mathbb{R}^n \to \mathbb{F}^n$ preserves unit distance, then f is injective.

Proof. We know that $D_n(\mathbb{F})$ is a dense subset of $(0, \infty)$. Therefore, for any $x, y \in \mathbb{R}^n$, $x \neq y$ there exists $z \in \mathbb{R}^n$ such that $|z - x| \neq |z - y|$ and $|z - x|, |z - y| \in D_n(\mathbb{F})$. All distances in $D_n(\mathbb{F})$ are preserved by f. Suppose $f(x) = f(y)$. This would imply that $|z - x|^2 = \varphi_n(z, x) = \varphi_n(f(z), f(x)) = \varphi_n(f(z), f(y)) = \varphi_n(z, y) = |z - y|^2$, which is a contradiction.

We shall prove that $\{d > 0 : d^2 \in \mathbb{Q}\} \subseteq D_2(\mathbb{F})$. We need the following technical Propositions 2-5.

Proposition 2 (cf. [5], [6]). The points $c_1 = (z_{1,1}, ..., z_{1,n}), ..., c_{n+1} = (z_{n+1,1}, ..., z_{n+1,n}) \in \mathbb{F}^n$ are affinely dependent if and only if their Cayley-Menger determinant $\Delta(c_1, ..., c_{n+1}) :=$

$$\det \begin{bmatrix} 0 & 1 & 1 & ... & 1 \\ 1 & 0 & \varphi_n(c_1, c_2) & ... & \varphi_n(c_1, c_{n+1}) \\ 1 & \varphi_n(c_2, c_1) & 0 & ... & \varphi_n(c_2, c_{n+1}) \\ ... & ... & ... & ... & ... \\ 1 & \varphi_n(c_{n+1}, c_1) & \varphi_n(c_{n+1}, c_2) & ... & 0 \end{bmatrix}$$

equals 0.

Proof. It follows from the equality

$$\left(\det\begin{bmatrix} z_{1,1} & z_{1,2} & ... & z_{1,n} & 1 \\ z_{2,1} & z_{2,2} & ... & z_{2,n} & 1 \\ ... & ... & ... & ... & ... \\ z_{n+1,1} & z_{n+1,2} & ... & z_{n+1,n} & 1 \end{bmatrix}\right)^2 = \frac{(-1)^{n+1}}{2^n} \cdot \Delta(c_1, ..., c_{n+1}).$$

Proposition 3 (cf. [5], [6]). For any points $c_1, ..., c_{n+k} \in \mathbb{F}^n$ $(k = 2, 3, 4, ...)$ their Cayley-Menger determinant equals 0 i.e. $\Delta(c_1, ..., c_{n+k}) = 0$.

Proof. Assume that $c_1 = (z_{1,1}, ..., z_{1,n}), ..., c_{n+k} = (z_{n+k,1}, ..., z_{n+k,n})$. The points $\widetilde{c}_1 = (z_{1,1}, ..., z_{1,n}, 0, ..., 0)$, $\widetilde{c}_2 = (z_{2,1}, ..., z_{2,n}, 0, ..., 0)$,..., $\widetilde{c}_{n+k} = (z_{n+k,1}, ..., z_{n+k,n}, 0, ..., 0) \in \mathbb{F}^{n+k-1}$ are affinely dependent. Since $\varphi_n(c_i, c_j) = \varphi_{n+k-1}(\widetilde{c}_i, \widetilde{c}_j)$ $(1 \le i \le j \le n + k)$ the Cayley-Menger determinant of points $c_1, ..., c_{n+k}$ is equal to the Cayley-Menger determinant of points $\widetilde{c}_1, ..., \widetilde{c}_{n+k}$ which equals 0 according to Proposition 2.

Proposition 4a. If $d \in \mathbb{F}$, $d \neq 0$, $c_1, ..., c_{n+1} \in \mathbb{F}^n$ and $\varphi_n(c_i, c_j) = d^2$ $(1 \le i < j \le n + 1)$, then the points c_1,...,c_{n+1} are affinely independent.

Proof. It follows from Proposition 2 because the Cayley-Menger determinant $\Delta(c_1, ..., c_{n+1}) =$

$$\det\begin{bmatrix} 0 & 1 & 1 & ... & 1 & 1 \\ 1 & 0 & d^2 & ... & d^2 & d^2 \\ 1 & d^2 & 0 & ... & d^2 & d^2 \\ ... & ... & ... & ... & ... & ... \\ 1 & d^2 & d^2 & ... & 0 & d^2 \\ 1 & d^2 & d^2 & ... & d^2 & 0 \end{bmatrix} = (-1)^{n+1}(n + 1)d^{2n} \neq 0.$$

Proposition 4b. If $a, b \in \mathbb{F}$, $a + b \neq 0$, $z, x, \widetilde{x} \in \mathbb{F}^2$ and $\varphi_2(z, x) = a^2$, $\varphi_2(x, \widetilde{x}) = b^2$, $\varphi_2(z, \widetilde{x}) = (a + b)^2$, then $\overrightarrow{zx} = \frac{a}{a+b}\overrightarrow{z\widetilde{x}}$.

Proof. Since $\varphi_2(z, \widetilde{x}) = (a + b)^2 \neq 0$ we conclude that $z \neq \widetilde{x}$. The Cayley-Menger determinant $\Delta(z, x, \widetilde{x}) =$

$$\det\begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & a^2 & (a+b)^2 \\ 1 & a^2 & 0 & b^2 \\ 1 & (a+b)^2 & b^2 & 0 \end{bmatrix} = 0,$$

so by Proposition 2 the points $z, x, \widetilde{x}$ are affinely dependent. Therefore, there exists $c \in \mathbb{F}$ such that $\overrightarrow{zx} = c \cdot \overrightarrow{z\widetilde{x}}$. Hence $a^2 = \varphi_2(z,x) = c^2 \cdot \varphi_2(z,\widetilde{x}) = c^2 \cdot (a+b)^2$. Thus $c = \frac{a}{a+b}$ or $c = -\frac{a}{a+b}$. If $c = -\frac{a}{a+b}$ then $\overrightarrow{x\widetilde{x}} = \overrightarrow{z\widetilde{x}} - \overrightarrow{zx} = \overrightarrow{z\widetilde{x}} + \frac{a}{a+b}\overrightarrow{z\widetilde{x}} = \frac{2a+b}{a+b}\overrightarrow{z\widetilde{x}}$. Hence $b^2 = \varphi_2(x,\widetilde{x}) = \left(\frac{2a+b}{a+b}\right)^2 \cdot \varphi_2(z,\widetilde{x}) = \left(\frac{2a+b}{a+b}\right)^2 \cdot (a+b)^2 = (2a+b)^2$. Therefore $0 = (2a+b)^2 - b^2 = 4a(a+b)$. Since $a+b \neq 0$ we conclude that $a = 0$, so $c = -\frac{a}{a+b} = \frac{a}{a+b}$ and the proof is completed.

Proposition 5 (cf. [6, Lemma, page 127] for $\mathbb{R}^n$). If x, y, c_0, c_1, $c_2 \in \mathbb{F}^2$, $\varphi_2(x,c_0) = \varphi_2(y,c_0)$, $\varphi_2(x,c_1) = \varphi_2(y,c_1)$, $\varphi_2(x,c_2) = \varphi_2(y,c_2)$ and c_0, c_1, c_2 are affinely independent, then $x = y$.

Proof. Computing we obtain that the vector $\overrightarrow{xy}$ is perpendicular to each of the linearly independent vectors $\overrightarrow{c_0c_1}$, $\overrightarrow{c_0c_2}$. Thus the vector $\overrightarrow{xy}$ is perpendicular to every linear combination of vectors $\overrightarrow{c_0c_1}$ and $\overrightarrow{c_0c_2}$. In particular, the vector $\overrightarrow{xy}$ is perpendicular to each of the vectors $[1,0]$, $[0,1]$. Therefore $\overrightarrow{xy} = 0$ and the proof is completed.

Lemma 1. If $d \in D_2(\mathbb{F})$ then $\sqrt{3} \cdot d \in D_2(\mathbb{F})$.

Proof. Let $d \in D_2(\mathbb{F})$, $x, y \in \mathbb{R}^2$ and $|x-y| = \sqrt{3} \cdot d$. Using the notation of Figure 1 we show that

$$S_{xy} := S_{y\widetilde{y}} \cup \bigcup_{i=1}^{2} S_{xp_i} \cup \bigcup_{i=1}^{2} S_{yp_i} \cup S_{p_1p_2} \cup \bigcup_{i=1}^{2} S_{x\widetilde{p_i}} \cup \bigcup_{i=1}^{2} S_{\widetilde{y}\widetilde{p_i}} \cup S_{\widetilde{p_1}\widetilde{p_2}}$$

satisfies condition $(*)$.



Figure 1

$$|x-y| = |x-\widetilde{y}| = \sqrt{3} \cdot d,\ |y-\widetilde{y}| = |p_1-p_2| = |\widetilde{p_1}-\widetilde{p_2}| = d$$
$$|x-p_i| = |y-p_i| = |x-\widetilde{p_i}| = |\widetilde{y}-\widetilde{p_i}| = d\ (i=1,2)$$

Assume that $f : S_{xy} \to \mathbb{F}^2$ preserves unit distance. Since

$$S_{xy} \supseteq S_{y\widetilde{y}} \cup \bigcup_{i=1}^{2} S_{xp_i} \cup \bigcup_{i=1}^{2} S_{yp_i} \cup S_{p_1p_2}$$

we conclude that f preserves the distances between y and $\widetilde{y}$, x and p_i $(i = 1, 2)$, y and p_i $(i = 1, 2)$, p_1 and p_2. Hence $\varphi_2(f(y), f(\widetilde{y})) = \varphi_2(f(x), f(p_i)) = \varphi_2(f(y), f(p_i)) = \varphi_2(f(p_1), f(p_2)) = d^2$ $(i = 1, 2)$. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x), f(p_1), f(p_2), f(y))$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & 0 & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & 0 & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & 0 \end{bmatrix} = 0.$$

Therefore

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 & t \\ 1 & d^2 & 0 & d^2 & d^2 \\ 1 & d^2 & d^2 & 0 & d^2 \\ 1 & t & d^2 & d^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$2d^2t \cdot (3d^2 - t) = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 3d^2 = |x - y|^2$$

or

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 0.$$

Analogously we may prove that

$$\varphi_2(f(x), f(\widetilde{y})) = \varphi_2(f(\widetilde{y}), f(x)) = 3d^2 = |x - \widetilde{y}|^2$$

or

$$\varphi_2(f(x), f(\widetilde{y})) = \varphi_2(f(\widetilde{y}), f(x)) = 0.$$

If $t = 0$ then the points $f(x)$ and $f(y)$ satisfy:

$$\begin{aligned}\varphi_2(f(x), f(x)) = 0 = \varphi_2(f(y), f(x)),\\ \varphi_2(f(x), f(p_1)) = d^2 = \varphi_2(f(y), f(p_1)),\\ \varphi_2(f(x), f(p_2)) = d^2 = \varphi_2(f(y), f(p_2)).\end{aligned}$$

By Proposition 4a the points $f(x)$, $f(p_1)$, $f(p_2)$ are affinely independent. Therefore by Proposition 5 $f(x) = f(y)$ and consequently

$$d^2 = \varphi_2(f(y), f(\widetilde{y})) = \varphi_2(f(x), f(\widetilde{y})) \in \{3d^2,\ 0\}.$$

Since $d^2 \neq 3d^2$ and $d^2 \neq 0$ we conclude that the case $t = 0$ cannot occur. This completes the proof of Lemma 1.

Lemma 2. If $d \in D_2(\mathbb{F})$ then $2 \cdot d \in D_2(\mathbb{F})$.

Proof. Let $d \in D_2(\mathbb{F})$, $x, y \in \mathbb{R}^2$ and $|x - y| = 2 \cdot d$. Using the notation of Figure 2 we show that

$$S_{xy} := \bigcup \{S_{ab} : a, b \in \{x, y, p_1, p_2, p_3\}, |a - b| = d \vee |a - b| = \sqrt{3} \cdot d\}$$

(where S_{xp_3} and S_{yp_2} are known to exist by Lemma 1) satisfies condition $(*)$.



Figure 2

$$|x - y| = 2 \cdot d$$
$$|p_1 - p_2| = |p_1 - p_3| = |p_2 - p_3| = |x - p_1| = |x - p_2| = |y - p_1| = |y - p_3| = d$$
$$|x - p_3| = |y - p_2| = \sqrt{3} \cdot d$$

Assume that $f : S_{xy} \to \mathbb{F}^2$ preserves unit distance. Then f preserves all distances between p_i and p_j $(1 \leq i < j \leq 3)$, x and p_i $(1 \leq i \leq 3)$, y and p_i $(1 \leq i \leq 3)$. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x),\ f(p_1),\ f(p_2),\ f(p_3),\ f(y))$ equals 0 i.e.

$$\det\begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & \varphi_2(f(x),f(p_1)) & \varphi_2(f(x),f(p_2)) & \varphi_2(f(x),f(p_3)) & \varphi_2(f(x),f(y)) \\ 1 & \varphi_2(f(p_1),f(x)) & 0 & \varphi_2(f(p_1),f(p_2)) & \varphi_2(f(p_1),f(p_3)) & \varphi_2(f(p_1),f(y)) \\ 1 & \varphi_2(f(p_2),f(x)) & \varphi_2(f(p_2),f(p_1)) & 0 & \varphi_2(f(p_2),f(p_3)) & \varphi_2(f(p_2),f(y)) \\ 1 & \varphi_2(f(p_3),f(x)) & \varphi_2(f(p_3),f(p_1)) & \varphi_2(f(p_3),f(p_2)) & 0 & \varphi_2(f(p_3),f(y)) \\ 1 & \varphi_2(f(y),f(x)) & \varphi_2(f(y),f(p_1)) & \varphi_2(f(y),f(p_2)) & \varphi_2(f(y),f(p_3)) & 0 \end{bmatrix} = 0.$$

Therefore

$$\det\begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & d^2 & d^2 & 3d^2 & t \\ 1 & d^2 & 0 & d^2 & d^2 & d^2 \\ 1 & d^2 & d^2 & 0 & d^2 & 3d^2 \\ 1 & 3d^2 & d^2 & d^2 & 0 & d^2 \\ 1 & t & d^2 & 3d^2 & d^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$3d^4 \cdot (t - 4d^2)^2 = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 4d^2 = |x-y|^2.$$

Lemma 3. If $a, b \in D_2(\mathbb{F})$ and $a > b$, then $\sqrt{a^2 - b^2} \in D_2(\mathbb{F})$.

Proof. Let $a, b \in D_2(\mathbb{F})$, $a > b$, $x, y \in \mathbb{R}^2$ and $|x - y| = \sqrt{a^2 - b^2}$. Using the notation of Figure 3 we show that

$$S_{xy} := S_{xp_1} \cup S_{xp_2} \cup S_{yp_1} \cup S_{yp_2} \cup S_{p_1p_2}$$

(where $S_{p_1p_2}$ is known to exist by Lemma 2) satisfies condition $(*)$.



Figure 3

$|x - y| = \sqrt{a^2 - b^2}$

$|x - p_1| = |x - p_2| = b$, $|y - p_1| = |y - p_2| = a$, $|p_1 - p_2| = 2b$

Assume that $f : S_{xy} \longrightarrow \mathbb{F}^2$ preserves unit distance. Then f preserves the distances between x and p_i $(i = 1, 2)$, y and p_i $(i = 1, 2)$, p_1 and p_2. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x),\ f(p_1),\ f(p_2),\ f(y))$ equals 0 i.e.

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & \varphi_2(f(x), f(p_1)) & \varphi_2(f(x), f(p_2)) & \varphi_2(f(x), f(y)) \\ 1 & \varphi_2(f(p_1), f(x)) & 0 & \varphi_2(f(p_1), f(p_2)) & \varphi_2(f(p_1), f(y)) \\ 1 & \varphi_2(f(p_2), f(x)) & \varphi_2(f(p_2), f(p_1)) & 0 & \varphi_2(f(p_2), f(y)) \\ 1 & \varphi_2(f(y), f(x)) & \varphi_2(f(y), f(p_1)) & \varphi_2(f(y), f(p_2)) & 0 \end{bmatrix} = 0.$$

Therefore

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & b^2 & b^2 & t \\ 1 & b^2 & 0 & 4b^2 & a^2 \\ 1 & b^2 & 4b^2 & 0 & a^2 \\ 1 & t & a^2 & a^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$-8b^2 \cdot (t + b^2 - a^2)^2 = 0.$$

Therefore

$$t = a^2 - b^2 = |x - y|^2.$$

Lemma 4. For each $n \in \{1, 2, 3, ...\}$ $\sqrt{n} \in D_2(\mathbb{F})$.

Proof. If $n \in \{2, 3, 4, ...\}$ and $\sqrt{n} \in D_2(\mathbb{F})$, then $\sqrt{n-1} = \sqrt{(\sqrt{n})^2 - 1^2} \in D_2(\mathbb{F})$; it follows from Lemma 3 and $1 \in D_2(\mathbb{F})$. On the other hand, by Lemma 2 all numbers $\sqrt{2^{2k}} = 2^k$ $(k = 0, 1, 2, ..)$ belong to $D_2(\mathbb{F})$. These two facts imply that all distances $\sqrt{n}$ $(n = 1, 2, 3, ...)$ belong to $D_2(\mathbb{F})$.

From Lemma 4 we obtain:

Lemma 5. For each $n \in \{1, 2, 3, ...\}$ $n = \sqrt{n^2} \in D_2(\mathbb{F})$.

Lemma 6. If $d \in D_2(\mathbb{F})$ then all distances $\dfrac{d}{k}$ $(k = 2, 3, 4, ...)$ belong to $D_2(\mathbb{F})$.

Proof. Let $d \in D_2(\mathbb{F})$, $k \in \{2, 3, 4, ...\}$, $x, y \in \mathbb{R}^2$ and $|x - y| = \dfrac{d}{k}$. We choose an integer $e \geq d$. Using the notation of Figure 4 we show that

$$S_{xy} := S_{\widetilde{x}\widetilde{y}} \cup S_{zx} \cup S_{x\widetilde{x}} \cup S_{z\widetilde{x}} \cup S_{zy} \cup S_{y\widetilde{y}} \cup S_{z\widetilde{y}}$$

(where sets S_{zx}, $S_{x\widetilde{x}}$, $S_{z\widetilde{x}}$, S_{zy}, $S_{y\widetilde{y}}$, $S_{z\widetilde{y}}$ corresponding to integer distances are known to exist by Lemma 5) satisfies condition $(*)$.



Figure 4

$|x-y| = \frac{d}{k}$

Assume that $f : S_{xy} \to \mathbb{F}^2$ preserves unit distance. Since $S_{xy} \supseteq S_{zx} \cup S_{x\widetilde{x}} \cup S_{z\widetilde{x}}$ we conclude that:

$$\varphi_2(f(z), f(x)) = \varphi_2(z, x) = e^2,$$
$$\varphi_2(f(x), f(\widetilde{x})) = \varphi_2(x, \widetilde{x}) = ((k-1)e)^2,$$
$$\varphi_2(f(z), f(\widetilde{x})) = \varphi_2(z, \widetilde{x}) = (ke)^2.$$

By Proposition 4b:

$$\overrightarrow{f(z)f(x)} = \frac{1}{k}\overrightarrow{f(z)f(\widetilde{x})} \qquad (1).$$

Analogously:

$$\overrightarrow{f(z)f(y)} = \frac{1}{k}\overrightarrow{f(z)f(\widetilde{y})} \qquad (2).$$

By (1) and (2):

$$\overrightarrow{f(x)f(y)} = \frac{1}{k}\overrightarrow{f(\widetilde{x})f(\widetilde{y})} \qquad (3).$$

Since $S_{xy} \supseteq S_{\widetilde{x}\widetilde{y}}$ we conclude that

$$\varphi_2(f(\widetilde{x}), f(\widetilde{y})) = \varphi_2(\widetilde{x}, \widetilde{y}) = d^2 \qquad (4).$$

By (3) and (4): $\varphi_2(f(x), f(y)) = \left(\frac{d}{k}\right)^2$ and the proof is completed.

Theorem 3. If $x, y \in \mathbb{R}^2$ and $|x-y|^2$ is a rational number, then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^2$ such that any map $f : S_{xy} \to \mathbb{F}^2$ that preserves unit distance preserves also the distance between x and y; in other words $\{d > 0 : d^2 \in \mathbb{Q}\} \subseteq D_2(\mathbb{F})$.

Proof. We need to prove that $\sqrt{\frac{p}{q}} \in D_2(\mathbb{F})$ for all positive integers p, q. Since $\sqrt{\frac{p}{q}} = \frac{\sqrt{pq}}{q}$ the assertion follows from Lemmas 4 and 6.

Conjecture 2. For each $n \geq 2$ $\{d > 0 : d^2 \in \mathbb{Q}\} \subseteq D_n(\mathbb{F})$.

As a corollary of Theorem 3 and $D_2(\mathbb{C}) \subseteq A_2(\mathbb{C}) \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$, we get:

Corollary 1. $D_2(\mathbb{C}) = A_2(\mathbb{C}) = \{d > 0 : d^2 \in \mathbb{Q}\}$.

The next theorems show which geometric properties are preserved by unit-distance preserving mappings.

Theorem 4. If $n \geq 2$, $f : \mathbb{R}^n \to \mathbb{F}^n$ preserves unit distance, $x, y \in \mathbb{R}^n$ and $x \neq y$, then $\varphi_n(f(x), f(y)) \neq 0$; in other words f preserves the relation that two points are at non-zero distance.

Proof. By Theorem 2 $f(x) \neq f(y)$. We know that $D_n(\mathbb{F})$ is a dense subset of $(0, \infty)$. In particular, $D_n(\mathbb{F})$ is unbounded from above. Therefore, there exist $d \in D_n(\mathbb{F})$ and $p_1, ..., p_{n-1}, p_n \in \mathbb{R}^n$ such that $|y - p_i| = d$ $(1 \leq i \leq n-1)$ and the points $x, p_1, ..., p_{n-1}, p_n$ are at distance d from each other. Since f preserves distance d, we conclude that $\varphi_n(f(y), f(p_i)) = d^2$ $(1 \leq i \leq n-1)$ and f preserves all distances between $x, p_1, ..., p_{n-1}, p_n$. Thus $\overrightarrow{f(x)f(p_i)} \cdot \overrightarrow{f(x)f(p_j)} = \frac{d^2}{2}$ $(1 \leq i < j \leq n)$ and the points $f(x), f(p_1), ..., f(p_{n-1}), f(p_n)$ are affinely independent by Proposition 4a. In particular, the vectors $\overrightarrow{f(x)f(p_i)}$ $(1 \leq i \leq n-1)$ are linearly independent. Assume, on the contrary, that $\varphi_n(f(x), f(y)) = \varphi_n(f(y), f(x)) = 0$. It implies that

$$\overrightarrow{f(x)f(y)} \text{ is perpendicular to each of the vectors } \overrightarrow{f(x)f(p_i)} \ (1 \leq i \leq n-1) \tag{5}$$

and the Cayley-Menger determinant $\Delta(f(x), f(p_1), ..., f(p_{n-1}), f(y)) =$

$$\det \begin{bmatrix} 0 & 1 & 1 & ... & 1 & 1 \\ \mathbf{1} & \mathbf{0} & \mathbf{d^2} & ... & \mathbf{d^2} & \mathbf{0} \\ 1 & d^2 & 0 & ... & d^2 & d^2 \\ ... & ... & ... & ... & ... & ... \\ 1 & d^2 & d^2 & ... & 0 & d^2 \\ \mathbf{1} & \mathbf{0} & \mathbf{d^2} & ... & \mathbf{d^2} & \mathbf{0} \end{bmatrix} = 0,$$

so by Proposition 2 the points $f(x), f(p_1), ..., f(p_{n-1}), f(y) \in \mathbb{F}^n$ are affinely dependent. Let

$$\overrightarrow{f(x)f(y)} = \alpha_1 \overrightarrow{f(x)f(p_1)} + ... + \alpha_{n-1} \overrightarrow{f(x)f(p_{n-1})} \tag{6}$$

where $\alpha_1, ..., \alpha_{n-1} \in \mathbb{F}$ and

$$\begin{aligned} \overrightarrow{f(x)f(y)} &= (t_1, ..., t_n) \in \mathbb{F}^n, \\ \overrightarrow{f(x)f(p_1)} &= (p_{1,1}, ..., p_{1,n}) \in \mathbb{F}^n, \\ &. \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \\ \overrightarrow{f(x)f(p_{n-1})} &= (p_{n-1,1}, ..., p_{n-1,n}) \in \mathbb{F}^n. \end{aligned}$$

By (5):

$$\begin{gathered} 0 = \overrightarrow{f(x)f(p_1)} \cdot \overrightarrow{f(x)f(y)} = p_{1,1} \cdot t_1 + ... + p_{1,n} \cdot t_n, \\ . \\ 0 = \overrightarrow{f(x)f(p_{n-1})} \cdot \overrightarrow{f(x)f(y)} = p_{n-1,1} \cdot t_1 + ... + p_{n-1,n} \cdot t_n. \end{gathered}$$

By (6):

$$\begin{gathered} t_1 = \alpha_1 \cdot p_{1,1} + ... + \alpha_{n-1} \cdot p_{n-1,1}, \\ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \\ t_n = \alpha_1 \cdot p_{1,n} + ... + \alpha_{n-1} \cdot p_{n-1,n}. \end{gathered}$$

Computing we obtain:

$$\begin{gathered} (\overrightarrow{f(x)f(p_1)} \cdot \overrightarrow{f(x)f(p_1)})\alpha_1 + ... + (\overrightarrow{f(x)f(p_1)} \cdot \overrightarrow{f(x)f(p_{n-1})})\alpha_{n-1} = 0, \\ . \\ (\overrightarrow{f(x)f(p_{n-1})} \cdot \overrightarrow{f(x)f(p_1)})\alpha_1 + ... + (\overrightarrow{f(x)f(p_{n-1})} \cdot \overrightarrow{f(x)f(p_{n-1})})\alpha_{n-1} = 0. \end{gathered}$$

Therefore:

$$\begin{gathered} d^2 \cdot \alpha_1 + \frac{d^2}{2} \cdot \alpha_2 + ... + \frac{d^2}{2} \cdot \alpha_{n-1} = 0, \\ . \\ \frac{d^2}{2} \cdot \alpha_1 + ... + \frac{d^2}{2} \cdot \alpha_{n-2} + d^2 \cdot \alpha_{n-1} = 0. \end{gathered}$$

Hence $\alpha_1 = ... = \alpha_{n-1} = 0$ and $\overrightarrow{f(x)f(y)} = 0$, which is a contradiction. The proof is completed.

At this point we consider only mappings from $\mathbb{R}^2$ to $\mathbb{F}^2$. We omit an easy proof of Proposition 6.

Proposition 6. If $E, F, C, D \in \mathbb{F}^2$, $\varphi_2(E,F) \neq 0$, $C \neq D$ and $\varphi_2(E,C) = \varphi_2(F,C) = \varphi_2(E,D) = \varphi_2(F,D)$, then $\overrightarrow{EC} = \overrightarrow{DF}$ and $\overrightarrow{FC} = \overrightarrow{DE}$ (see the points E, F, C and D in Figure 5).

Theorem 5. If $t \in \mathbb{Q}$, $0 < t < 1$, $A, B \in \mathbb{R}^2$, $A \neq B$, $C = tA + (1-t)B$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $f(C) = tf(A) + (1-t)f(B)$.

Proof. We choose $r \in \mathbb{Q}$ such that $r > |AB|$ and $r < \frac{|AB|}{|1-2t|}$ if $t \neq \frac{1}{2}$. There exists $D \in \mathbb{R}^2$ such that $|AD| = (1-t)r$ and $|BD| = tr$. Let $E = tA + (1-t)D$ and $F = (1-t)B + tD$, see Figure 5.



Figure 5

The segments AE, ED, AD, BF, FD, BD, EC and FC have rational lengths and $|EC| = |FC| = |ED| = |FD| = t(1-t)r$. By Theorem 3 f preserves rational distances. Therefore:

$$\varphi_2(f(A), f(E)) = \varphi_2(A, E) = |AE|^2 = ((1-t)^2 r)^2,$$
$$\varphi_2(f(E), f(D)) = \varphi_2(E, D) = |ED|^2 = (t(1-t)r)^2,$$
$$\varphi_2(f(A), f(D)) = \varphi_2(A, D) = |AD|^2 = ((1-t)r)^2.$$

Since $(1-t)^2 r + t(1-t)r = (1-t)r$, by Proposition 4b $f(E) = tf(A) + (1-t)f(D)$. Analogously $f(F) = (1-t)f(B) + tf(D)$. Since $C \neq D$, by Theorem 2 $f(C) \neq f(D)$. Since $E \neq F$, by Theorem 4 $\varphi_2(f(E), f(F)) \neq 0$. By Theorem 3 f preserves rational distances. Therefore:

$$\varphi_2(f(E), f(C)) = \varphi_2(E, C) = |EC|^2 = (t(1-t)r)^2,$$
$$\varphi_2(f(F), f(C)) = \varphi_2(F, C) = |FC|^2 = (t(1-t)r)^2,$$
$$\varphi_2(f(E), f(D)) = \varphi_2(E, D) = |ED|^2 = (t(1-t)r)^2,$$
$$\varphi_2(f(F), f(D)) = \varphi_2(F, D) = |FD|^2 = (t(1-t)r)^2.$$

By Proposition 6 $\overrightarrow{f(E)F(C)} = \overrightarrow{f(D)f(F)}$, so $\overrightarrow{f(A)f(C)} = \overrightarrow{f(A)f(E)} + \overrightarrow{f(E)f(C)} = \overrightarrow{f(A)f(E)} + \overrightarrow{f(D)f(F)} = (1-t)\overrightarrow{f(A)f(D)} + (1-t)\overrightarrow{f(D)f(B)} = (1-t)\overrightarrow{f(A)f(B)}$ and the proof is completed.

It is easy to show that the present form of Theorem 5 implies a more general form without the assumptions $0 < t < 1$ and $A \neq B$.

Lemma 7. If $A, B, C, D \in \mathbb{R}^2$, $\overrightarrow{AB} = \overrightarrow{CD}$, $|AC| = |BD| \in \mathbb{Q}$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\overrightarrow{f(A)f(B)} = \overrightarrow{f(C)f(D)}$.

Proof. There exist $m \in \{0, 1, 2, ...\}$ and $A_0, C_0, A_1, C_1, ..., A_m, C_m \in \mathbb{R}^2$ such that $A_0 = A$, $C_0 = C$, $A_m = B$, $C_m = D$ and for each $i \in \{0, 1, ..., m-1\}$ $A_iC_iC_{i+1}A_{i+1}$ is a rhombus with a rational side, see Figure 6 where $m = 3$.



Figure 6

By Theorem 3 for each $i \in \{0, 1, ..., m-1\}$ f preserves the lenghts of the sides of the rhombus $A_iC_iC_{i+1}A_{i+1}$. For each $i \in \{0, 1, ..., m-1\}$ we have: $\varphi_2(f(A_i), f(C_{i+1})) \neq 0$ (by Theorem 4) and $f(C_i) \neq f(A_{i+1})$ (by Theorem 2). Therefore, by Proposition 6 for each $i \in \{0, 1, ..., m-1\}$ $\overrightarrow{f(A_i)f(A_{i+1})} = \overrightarrow{f(C_i)f(C_{i+1})}$. Hence $\overrightarrow{f(A)f(B)} = \overrightarrow{f(A_0)f(A_m)} = \overrightarrow{f(A_0)f(A_1)} + \overrightarrow{f(A_1)f(A_2)} + ... + \overrightarrow{f(A_{m-1})f(A_m)} = \overrightarrow{f(C_0)f(C_1)} + \overrightarrow{f(C_1)f(C_2)} + ... + \overrightarrow{f(C_{m-1})f(C_m)} = \overrightarrow{f(C)f(D)}$.

Theorem 6. If $A, B, C, D \in \mathbb{R}^2$, $\overrightarrow{AB} = \overrightarrow{CD}$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\overrightarrow{f(A)f(B)} = \overrightarrow{f(C)f(D)}$.

Proof. There exist $E, F \in \mathbb{R}^2$ such that $\overrightarrow{AB} = \overrightarrow{EF} = \overrightarrow{CD}$, $|AE| = |BF| \in \mathbb{Q}$ and $|EC| = |FD| \in \mathbb{Q}$, see Figure 7.



Figure 7

By Lemma 7 $\overrightarrow{f(A)f(B)} = \overrightarrow{f(E)f(F)}$ and $\overrightarrow{f(E)f(F)} = \overrightarrow{f(C)f(D)}$, so $\overrightarrow{f(A)f(B)} = \overrightarrow{f(C)f(D)}$.

As a corollary of Theorems 5 and 6 we get:

Theorem 7. If $A, B, C, D \in \mathbb{R}^2$, $r \in \mathbb{Q}$, $\overrightarrow{CD} = r\overrightarrow{AB}$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\overrightarrow{f(C)f(D)} = r\overrightarrow{f(A)f(B)}$.

Lemma 8. If $P, Q, X, Y \in \mathbb{R}^2$, $\overrightarrow{PQ}$ is perpendicular to $\overrightarrow{XY}$, $|XY| \in \mathbb{Q}$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\overrightarrow{f(P)f(Q)}$ is perpendicular to $\overrightarrow{f(X)f(Y)}$.

Proof. There exist $A, B, C, D, E, F \in \mathbb{R}^2$ and $r, s \in \mathbb{Q}$ such that $\overrightarrow{PQ} = r\overrightarrow{DE}$, $\overrightarrow{XY} = s\overrightarrow{AB}$ and the points A, B, C, D, E, F form the configuration from Figure 8; it is a part of Kempe's linkage for drawing straight lines, see [12].



Figure 8

$$B \neq D, \;\; B \neq E$$

$$|AB| = |AD| = 4, \;\; |CB| = |CD| = |CE| = 2, \;\; |AF| = 3, \;\; |FB| = |FE| = 1$$

By Theorem 7 $\overrightarrow{f(P)f(Q)} = r\overrightarrow{f(D)f(E)}$ and $\overrightarrow{f(X)f(Y)} = s\overrightarrow{f(A)f(B)}$, so it suffices to prove that $\overrightarrow{f(D)f(E)} \cdot \overrightarrow{f(A)f(B)} = 0$. Let $a = \varphi_2(f(B), f(D))$, $b = \varphi_2(f(A), f(C))$, $c = \varphi_2(f(B), f(E))$, $d = \varphi_2(f(C), f(F))$, $e = \varphi_2(f(A), f(E))$.

Computing the value of c we obtain:
$c = \varphi_2(f(B), f(E)) = (\overrightarrow{f(B)f(E)})^2 = (\overrightarrow{f(A)f(E)} - \overrightarrow{f(A)f(B)})^2 =$
$e - 2\overrightarrow{f(A)f(E)} \cdot \overrightarrow{f(A)f(B)} + 16$, so

$$\overrightarrow{f(A)f(E)} \cdot \overrightarrow{f(A)f(B)} = 8 + \tfrac{1}{2}(e - c) \qquad (7).$$

Computing the value of a we obtain:
$a = \varphi_2(f(B), f(D)) = (\overrightarrow{f(B)f(D)})^2 = (\overrightarrow{f(A)f(D)} - \overrightarrow{f(A)f(B)})^2 =$
$16 - 2\overrightarrow{f(A)f(D)} \cdot \overrightarrow{f(A)f(B)} + 16$, so

$$\overrightarrow{f(A)f(D)} \cdot \overrightarrow{f(A)f(B)} = 16 - \tfrac{1}{2}a \qquad (8).$$

The next calculations are based on Proposition 3 and the observation that distances 1, 2, 3 and 4 are preserved by f.

$$0 = \Delta(f(A), f(B), f(E), f(F)) = \det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 16 & e & 9 \\ 1 & 16 & 0 & c & 1 \\ 1 & e & c & 0 & 1 \\ 1 & 9 & 1 & 1 & 0 \end{bmatrix} = -2(e - 16 + 3c)^2, \text{ so}$$

$$e = 16 - 3c \qquad (9).$$

$$0 = \Delta(f(A), f(B), f(C), f(F)) = \det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 16 & b & 9 \\ 1 & 16 & 0 & 4 & 1 \\ 1 & b & 4 & 0 & d \\ 1 & 9 & 1 & d & 0 \end{bmatrix} = -2(b - 4d)^2, \text{ so } b = 4d.$$

Thus $0 = \Delta(f(A), f(B), f(C), f(D)) = \det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 16 & b & 16 \\ 1 & 16 & 0 & 4 & a \\ 1 & b & 4 & 0 & 4 \\ 1 & 16 & a & 4 & 0 \end{bmatrix} =$

$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 16 & 4d & 16 \\ 1 & 16 & 0 & 4 & a \\ 1 & 4d & 4 & 0 & 4 \\ 1 & 16 & a & 4 & 0 \end{bmatrix} = -8a(ad + 4(d^2 - 10d + 9))$. By Theorem 4 $a \neq 0$, so $a = -4\frac{d^2-10d+9}{d}$.

$0 = \Delta(f(B), f(C), f(E), f(F)) = \det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 4 & c & 1 \\ 1 & 4 & 0 & 4 & d \\ 1 & c & 4 & 0 & 1 \\ 1 & 1 & d & 1 & 0 \end{bmatrix} = -2c(cd + d^2 - 10d + 9)$. By Theorem 4 $c \neq 0$, so $c = -\frac{d^2-10d+9}{d}$. Therefore

$$a = 4c \qquad (10).$$

By (7)-(10):

$\overrightarrow{f(D)f(E)} \cdot \overrightarrow{f(A)f(B)} = (\overrightarrow{f(A)f(E)} - \overrightarrow{f(A)f(D)}) \cdot \overrightarrow{f(A)f(B)} =$
$\overrightarrow{f(A)f(E)} \cdot \overrightarrow{f(A)f(B)} - \overrightarrow{f(A)f(D)} \cdot \overrightarrow{f(A)f(B)} = 8 + \frac{1}{2}(e-c) - (16 - \frac{1}{2}a) =$
$\frac{1}{2}a - \frac{1}{2}c + \frac{1}{2}e - 8 = \frac{1}{2}a - \frac{1}{2}c + \frac{1}{2}(16-3c) - 8 = \frac{1}{2}a - 2c = 0.$

The proof is completed.

Theorem 8. If $A,B,C,D \in \mathbb{R}^2$, $\overrightarrow{AB}$ and $\overrightarrow{CD}$ are linearly dependent and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\overrightarrow{f(A)f(B)}$ and $\overrightarrow{f(C)f(D)}$ are linearly dependent.

Proof. We choose $X,Y \in \mathbb{R}^2$ such that $|XY| = 1$ and both vectors $\overrightarrow{AB}$ and $\overrightarrow{CD}$ are perpendicular to $\overrightarrow{XY}$. Then $\varphi_2(f(X),f(Y)) = 1$, so obviously $f(X) \neq f(Y)$. By Lemma 8 both vectors $\overrightarrow{f(A)f(B)}$ and $\overrightarrow{f(C)f(D)}$ are perpendicular to $\overrightarrow{f(X)f(Y)}$. These two facts imply that $\overrightarrow{f(A)f(B)}$ and $\overrightarrow{f(C)f(D)}$ are linearly dependent.

As a corollary of Theorem 8 we get:

Corollary 2. Unit-distance preserving mappings from $\mathbb{R}^2$ to $\mathbb{F}^2$ preserve collinearity of points.

Theorem 9. If $P,Q,X,Y \in \mathbb{R}^2$, $\overrightarrow{PQ}$ is perpendicular to $\overrightarrow{XY}$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\overrightarrow{f(P)f(Q)}$ is perpendicular to $\overrightarrow{f(X)f(Y)}$.

Proof. We may assume that $X \neq Y$. Set $Z = X + \frac{\overrightarrow{XY}}{|XY|}$. By Lemma 8 $\overrightarrow{f(P)f(Q)}$ and $\overrightarrow{f(X)f(Z)}$ are perpendicular. The points X, Y and Z are collinear, so by Corollary 2 $f(X)$, $f(Y)$ and $f(Z)$ are collinear. Since $|XZ| = 1$, we conclude that $\varphi_2(f(X),f(Z)) = 1$, so $f(X) \neq f(Z)$. Thus there exists $\alpha \in \mathbb{F}$ such that $\overrightarrow{f(X)f(Y)} = \alpha\overrightarrow{f(X)f(Z)}$. Hence $\overrightarrow{f(P)f(Q)} \cdot \overrightarrow{f(X)f(Y)} = \overrightarrow{f(P)f(Q)} \cdot \alpha\overrightarrow{f(X)f(Z)} = \alpha\left(\overrightarrow{f(P)f(Q)} \cdot \overrightarrow{f(X)f(Z)}\right) = \alpha \cdot 0 = 0$.

Lemma 9. If $A,X,Y \in \mathbb{R}^2$, $|AX| = |AY|$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\varphi_2(f(A),f(X)) = \varphi_2(f(A),f(Y))$.

Proof. Let $Z = \frac{1}{2}X + \frac{1}{2}Y$. By Theorem 5 $f(Z) = \frac{1}{2}f(X) + \frac{1}{2}f(Y)$. By Theorem 9 $\overrightarrow{f(A)f(Z)}$ is perpendicular to $\overrightarrow{f(Z)f(X)}$ and $\overrightarrow{f(A)f(Z)}$ is perpendicular to $\overrightarrow{f(Z)f(Y)}$. Hence $\varphi_2(f(A),f(X)) = \varphi_2(f(A),f(Z)) + \frac{1}{4}\varphi_2(f(X),f(Y)) = \varphi_2(f(A),f(Y))$.

Theorem 10. If $A,B,C,D \in \mathbb{R}^2$, $|AB| = |CD|$ and $f : \mathbb{R}^2 \to \mathbb{F}^2$ preserves unit distance, then $\varphi_2(f(A),f(B)) = \varphi_2(f(C),f(D))$.

Proof. We may assume that $|AB| = |CD| > 0$. There exist $m \in \{0, 1, 2, ...\}$ and points $T_0, T_1, ..., T_m \in \mathbb{R}^2$ such that $|AB| = |BT_0| = |T_0T_1| = ... = |T_{m-1}T_m| = |T_mC| = |CD|$. By Lemma 9 $\varphi_2(f(A), f(B)) = \varphi_2(f(B), f(T_0)) = \varphi_2(f(T_0), f(T_1)) = ... = \varphi_2(f(T_{m-1}), f(T_m)) = \varphi_2(f(T_m), f(C)) = \varphi_2(f(C), f(D))$.

We have already proved the main results that are needed for proving Conjecture 1 for $n = 2$. Such a proof will appear in the next publication.

Acknowledgement. The author wishes to thank Professor Mowaffaq Hajja for improvement of the proof of Proposition 1.

References

[1] F. S. Beckman and D. A. Quarles Jr., *On isometries of euclidean spaces*, Proc. Amer. Math. Soc. 4 (1953), 810–815.

[2] W. Benz, *An elementary proof of the theorem of Beckman and Quarles*, Elem. Math. 42 (1987), 4–9.

[3] W. Benz, *Geometrische Transformationen (unter besonderer Berücksichtigung der Lorentztransformationen)*, BI Wissenschaftsverlag, Mannheim, 1992.

[4] W. Benz, *Real geometries*, BI Wissenschaftsverlag, Mannheim, 1994.

[5] L. M. Blumenthal, *Theory and applications of distance geometry*, 2nd ed., Chelsea Publishing Co., New York, 1970.

[6] K. Borsuk, *Multidimensional analytic geometry*, Polish Scientific Publishers, Warsaw, 1969.

[7] N. Bourbaki, *Elements of mathematics, Algebra II, Chapters 4–7*, Springer-Verlag, Berlin, 1990.

[8] U. Everling, *Solution of the isometry problem stated by K. Ciesielski*, Math. Intelligencer 10 (1988), No. 4, page 47.

[9] D. J. H. Garling, *A course in Galois theory*, Cambridge Univ. Press, Cambridge, 1995.

[10] H. Kestelman, *Automorphisms of the field of complex numbers*, Proc. London Math. Soc., II. Ser. 53 (1951), 1–12.

[11] M. Kuczma, *An introduction to the theory of functional equations and inequalities: Cauchy's equation and Jensen's inequality*, Polish Scientific Publishers and Silesian University, Warsaw-Cracow-Katowice, 1985.

[12] H. Rademacher and O. Toeplitz, *The enjoyment of mathematics*, Princeton University Press, Princeton, 1994.

[13] A. Tyszka, *Discrete versions of the Beckman-Quarles theorem*, Aequationes Math. 59 (2000), 124–133.

[14] A. Tyszka, *Discrete versions of the Beckman-Quarles theorem from the definability results of R. M. Robinson*, Algebra, Geometry & their Applications, Seminar Proceedings 1 (2001), 88–90, Yerevan State University Press.

[15] A. Tyszka, *Beckman-Quarles type theorems for mappings from $\mathbb{R}^n$ to $\mathbb{C}^n$*, Aequationes Math., to appear.

[16] P. B. Yale, *Automorphisms of the complex numbers*, Math. Mag. 39 (1966), 135–141.

Apoloniusz Tyszka
Technical Faculty
Hugo Kołłątaj University
Balicka 104, 30-149 Kraków, Poland
E-mail: *rttyszka@cyf-kr.edu.pl*